

Example 2 We remind that

- A $K3$ surface is a smooth minimal algebraic surface S with $g(S) = 0$ and $h^1(S) = 0$.

$K3$ surfaces have invariants $h^2 = 0, g = 0, p_g = 1$
 $e = 24, h^1(S) = 0$.

- An Enriques surface is a smooth minimal algebraic surface S with $g(S) = 0$ and a non-trivial canonical bundle for which $2K_S = 0$.

Enriques surfaces have invariants $h^2_S = g = p_g = 0$
 $e = 12, h^1(S) = 0$

Historical Remark:

In 1895 Castelnuovo posed a question whether a surface with $g = p_g = 0$ is necessarily rational.

Enriques answered to this question: there are examples of surfaces with $g = p_g = 0$ that are not rational. These were the first examples of Enriques Surfaces.

Later on, Castelnuovo proved the beautiful Theorem (Castelnuovo Rationality Criterion)

S is a rational surface $\iff g(S) = 0$ and $P_2 = h^0(2K_S) = 0$.

We observe that any Enriques Surface is the quotient of an étale double covering of a K3 surface.

Indeed, given an Enriques Surface Y , then $2K_Y \equiv 0$, so $L_{\bar{1}} := K_Y$ and $D_{\bar{1}} = 0$ defines an abelian cover $\pi: X \rightarrow Y$ with group $G = \mathbb{Z}/2$. By construction, π is étale (no verification), so X is smooth minimal and by Riemann-Roch: $K_X = \pi^*(K_Y)$

Thus, $2K_X = \pi^*(2K_Y) = 0$, and since $L_{\bar{1}} = K_Y$, then $p_g(X) = h^0(X, K_X) = h^2(X, \mathcal{O}_X) = h^2(Y, \mathcal{O}_Y) + h^2(Y, L_{\bar{1}}^{-1}) = h^0(Y, K_Y) + h^0(Y, K_Y + L_{\bar{1}}) = 0 + 1 = 1$

\downarrow
 Serre Duality

So $2K_X \equiv 0$, $p_g(X) = 1 \Rightarrow K_X \equiv 0$.

Instead, $q(X) = h^1(X, \mathcal{O}_X) = h^1(Y, \mathcal{O}_Y) + h^1(Y, L_{\bar{1}}^{-1}) = 2h^1(Y, \mathcal{O}_Y) = 2q(Y) = 0$

$-K_Y = K_Y$

Thus, X is a K3 surface.

Remark K3 surfaces are always simply connected, so the universal cover of an Enriques surface is always a K3 surface.

WARNING: The opposite is false, namely is not

true that any $K3$ -surface has an involution $\sigma \in \text{Aut}(K3)$ s.t. $K3/\langle \sigma \rangle$ is Enriques.

Lemma 1 (the divisors D_g determine L_X up to torsion).
(F.-Pignatelli 2023, Cor. 1.6, Pord., Prop. 2.1)

Any invertible sheaf $L_X, X \in G^*$, satisfies

$$|X| \cdot L_X \equiv \sum_{g \in G} \left\lfloor \frac{|X| \cdot r_X^g}{|g|} \right\rfloor D_g$$

it denotes the order of X

proof We prove by Induction on k that

$$k \cdot L_X \equiv L_{X^k} + \sum_{g \in G} \left\lfloor \frac{k r_X^g}{|g|} \right\rfloor D_g$$

$k=1$ trivial, so suppose the thesis holds for $k-1$.

We use Pardini Equation: $L_X + L_{X^{k-1}} \equiv L_{X^k} + \sum_{r_X^g + r_{X^{k-1}}^g \geq |g|} D_g$

However, by inductive hypothesis, we have:

$$L_X + (k-1)L_X - \sum_{g \in G} \left\lfloor \frac{(k-1)r_X^g}{|g|} \right\rfloor D_g \equiv L_{X^k} + \sum_{r_X^g + r_{X^{k-1}}^g \geq |g|} D_g$$

$$\Rightarrow kL_X \equiv L_{X^k} + \sum_{g \in G} \left\lfloor \frac{(k-1)r_X^g}{|g|} \right\rfloor D_g + \sum_{r_X^g + r_{X^{k-1}}^g \geq |g|} D_g \quad (*)$$

Fixed $g \in G$, $(k-1)r_X^g = \alpha|g| + r_{X^{k-1}}^g$, where $\alpha = \left\lfloor \frac{(k-1)r_X^g}{|g|} \right\rfloor$ by definition.

Then $r_X^g + r_{X^{k-1}}^g = r_X^g + (k-1)r_X^g - \alpha|g| = kr_X^g - \alpha|g| \geq |g| \Leftrightarrow kr_X^g \geq (\alpha+1)|g| \Leftrightarrow \left\lfloor \frac{kr_X^g}{|g|} \right\rfloor = \alpha+1$
Instead, if $kr_X^g < (\alpha+1)|g| \Rightarrow \left\lfloor \frac{kr_X^g}{|g|} \right\rfloor = \alpha$

In both cases, D_g occurs in (*) with multiplicity $\left\lfloor \frac{kr_X^g}{|g|} \right\rfloor$.

Finally, for $k = |X|$, then $L_{\chi^k} = L_{\mathbb{1}} = 0$, and $\frac{|X| r_{\chi^k}}{|g|}$ is an integer because $\chi(g) = e^{\frac{2\pi i}{|g|} r_{\chi} g}$

$$\Rightarrow \chi^{|X|}(g) = 1 = e^{2\pi i \cdot \frac{|X| r_{\chi^k}}{|g|}}$$


As we have seen in Example 1 above, verifying that a set of building data determine an abelian cover can become computational challenging as the number of equations to satisfy grows quadratically with the order of G . The next proposition reduces the number of linear equations to satisfy to just the length of the group G .

Def (reduced building data) Let $\pi: X \rightarrow Y$ be a G -cover, Y smooth, X normal, with build. data $\{L_X, \{D_g\}$. Assume $G^* = \bigoplus_{i=1}^r \langle \chi_i \rangle$. Then $\{L_{\chi_i}, \{D_g\}_{g \in G}$ is called a set of reduced building data of π .

Proposition (Pardini, Prop. 2.1)

Let Y be a smooth algebraic variety and let G be an abelian group.

Let X_1, \dots, X_s be some generators of G^* , so $G^* = \langle X_1, \dots, X_s \rangle$.

Given an abelian cover $\pi: X \rightarrow Y$ with group G and building data $\{L_{X_i}\}_{X_i \in G^*}, \{D_g\}_{g \in G}$, then

$$\forall i=1, \dots, s, \quad |X_i| L_{X_i} \equiv \sum_{g \in G} \frac{|X_i| r_{X_i}^g}{|g|} D_g. \quad (\star)$$

Conversely, given a collection $\{L_{X_i}, \dots, L_{X_s}\}$ of line bundles of Y and divisors $\{D_g\}_{g \in G}$ such that (\star) hold for any X_1, \dots, X_s , then it there exists a abelian cover $\pi: X \rightarrow Y$ with group G .

When X is normal, then π has reduced building data $\{L_{X_i}, \dots, L_{X_s}\}, \{D_g\}_{g \in G}$.

If Y is complete, then the reduced building data determine the cover up to isomorph. of Galois covers.

proof Given an abelian cover π , then (\star) hold by the previous Lemma 1.

Furthermore, $\{L_{X_i}\}_{X_i \in G^*}$ are determined by $\{L_{X_1}, \dots, L_{X_s}\}$ and $\{D_g\}_{g \in G}$. To see this, we remind that:

1) $r_{y^s}^{\delta}$ is the remainder of $s \cdot r_y^{\delta}$ by the division with $|g|$

2) $\lfloor \frac{a+b}{c} \rfloor = \lfloor \frac{a}{c} \rfloor + \lfloor \frac{b}{c} \rfloor + \lfloor \frac{r_1+r_2}{c} \rfloor$, where $a = \alpha_1 c + r_1$
 $b = \alpha_2 c + r_2$

Thus, let us consider $x = x^{a_1} \dots x^{a_s} \in G^*$.

We need to write L_x in function of $\{x_i, i=1, \dots, s\}, \{D_g\}$.

$$L_{x_1^{a_1} \dots x_{s-1}^{a_{s-1}} x_s^{a_s}} = L_x + \sum_{g \in G} \left\lfloor \frac{r_{x_1^{a_1} \dots x_{s-1}^{a_{s-1}}}^{\delta} + r_{x_s^{a_s}}^{\delta}}{|g|} \right\rfloor D_g$$

$$\Rightarrow L_x = \underset{\substack{\downarrow \\ \text{by Lemma 1}}}{L_{x_1^{a_1} \dots x_{s-1}^{a_{s-1}}}} + a_s L_{x_s} - \sum_{g \in G} \left\lfloor \frac{a_s r_{x_s}^{\delta}}{|g|} \right\rfloor D_g - \sum_{g \in G} \left\lfloor \frac{r_{x_1^{a_1} \dots x_{s-1}^{a_{s-1}}}^{\delta} + r_{x_s^{a_s}}^{\delta}}{|g|} \right\rfloor D_g$$

$$= L_{x_1^{a_1} \dots x_{s-1}^{a_{s-1}}} + a_s L_{x_s} - \sum_{g \in G} \left\lfloor \frac{r_{x_1^{a_1} \dots x_{s-1}^{a_{s-1}}}^{\delta} + a_s r_{x_s}^{\delta}}{|g|} \right\rfloor D_g$$

$a_s r_{x_s}^{\delta} = a |g| + r_{x_s}^{\delta}$
 and by 2) we have

$$\left\lfloor \frac{r_{x_1^{a_1} \dots x_{s-1}^{a_{s-1}}}^{\delta} + a_s r_{x_s}^{\delta}}{|g|} \right\rfloor = 0 + \left\lfloor \frac{a_s r_{x_s}^{\delta}}{|g|} \right\rfloor + \left\lfloor \frac{r_{x_1^{a_1} \dots x_{s-1}^{a_{s-1}}}^{\delta} + r_{x_s}^{\delta}}{|g|} \right\rfloor$$

This computation suggests that

$$L_x = \sum_{i=1}^s a_i L_{x_i} - \sum_{g \in G} q_x^g D_g, \text{ where } q_x^g := \left\lfloor \frac{\sum_{i=1}^s a_i r_{x_i}^{\delta}}{|g|} \right\rfloor$$

Let us prove it by induction on s :

$s=1$: This is Lemma 1.

By the previous computation, we have

$$L_x = L_{x_1^{a_1} \dots x_{s-1}^{a_{s-1}}} + a_s L_{x_s} - \sum_{g \in G} \left\lfloor \frac{r_{x_1^{a_1} \dots x_{s-1}^{a_{s-1}}}^{\delta} + a_s r_{x_s}^{\delta}}{|g|} \right\rfloor D_g$$

$$= \sum_{i=1}^{s-1} a_i L_{x_i} - \sum_{g \in G} \left(\left\lfloor \frac{\sum_{i=1}^{s-1} a_i r_{x_i}^{\delta}}{|g|} \right\rfloor + \left\lfloor \frac{r_{x_1^{a_1} \dots x_{s-1}^{a_{s-1}}}^{\delta} + a_s r_{x_s}^{\delta}}{|g|} \right\rfloor \right) D_g$$

inductive hypothesis

$$\left\lfloor \frac{\sum_{i=1}^{s-1} a_i r_{x_i}^{\delta}}{|g|} \right\rfloor = \left\lfloor \frac{\sum_{i=1}^{s-1} a_i r_{x_i}^{\delta}}{|g|} \right\rfloor + \left\lfloor \frac{a_s r_{x_s}^{\delta}}{|g|} \right\rfloor + \left\lfloor \frac{r_{x_1^{a_1} \dots x_{s-1}^{a_{s-1}}}^{\delta} + r_{x_s}^{\delta}}{|g|} \right\rfloor = \left\lfloor \frac{\sum_{i=1}^{s-1} a_i r_{x_i}^{\delta}}{|g|} \right\rfloor + \left\lfloor \frac{r_{x_1^{a_1} \dots x_{s-1}^{a_{s-1}}}^{\delta} + a_s r_{x_s}^{\delta}}{|g|} \right\rfloor$$

$$= \sum_{i=1}^n a_i L_{x_i} - \sum_{g \in G} q_x^g D_g \text{ and the thesis follows.}$$

Conversely, let us consider a set of data $\{L_{x_i}\}_{i=1}^n$, $\{D_g\}_g$ of \mathcal{Y} satisfying (\star) . To prove the existence of a covering π , It is sufficient to define the remain L_x , $x \in G^*$, such that $\{L_x\}_{x \in G^*}$ and $\{D_g\}_g$ satisfy Pardini Equations.

The previous computation suggests how to define L_x :

$$L_x := \sum_{i=1}^n a_i L_{x_i} - \sum_{g \in G} q_x^g D_g, \text{ where } q_x^g := \left\lfloor \frac{\sum_{i=1}^n a_i r_{x_i}^g}{|g|} \right\rfloor$$

Let us prove $\{L_x\}_{x \in G^*}$ and $\{D_g\}_{g \in G}$ satisfy Pardini Equations:

$$\begin{aligned} L_x + L_\eta &= \sum_{i=1}^n a_i L_{x_i} - \sum_{g \in G} q_x^g D_g + \sum_{i=1}^n b_i L_{x_i} - \sum_{g \in G} q_\eta^g D_g \\ &\quad \begin{array}{l} x = x_1^{a_1} \dots x_s^{a_s} \\ \eta = x_1^{b_1} \dots x_s^{b_s} \end{array} \\ &= \sum_{i=1}^n (a_i + b_i) L_{x_i} - \sum_{g \in G} (q_x^g + q_\eta^g) D_g \\ &= L_{x\eta} + \sum_{g \in G} \left\lfloor \frac{r_x^g + r_\eta^g}{|g|} \right\rfloor D_g \text{ and we have done.} \end{aligned}$$

if we prove $q_{x\eta}^g = q_x^g + q_\eta^g - \left\lfloor \frac{r_x^g + r_\eta^g}{|g|} \right\rfloor$

$$q_x^g + q_\eta^g = \left\lfloor \frac{\sum_{i=1}^n a_i r_{x_i}^g}{|g|} \right\rfloor + \left\lfloor \frac{\sum_{i=1}^n b_i r_{x_i}^g}{|g|} \right\rfloor = \left\lfloor \frac{\sum a_i r_{x_i}^g}{|g|} \right\rfloor + \left\lfloor \frac{\sum b_i r_{x_i}^g}{|g|} \right\rfloor - \left\lfloor \frac{r_x^g + r_\eta^g}{|g|} \right\rfloor$$

by 2)



Example 3 If Y is simply connected, then $\text{Pic}(Y)$ has not torsion, i.e.
 $\forall D \in \text{Pic}(Y) \quad nD = 0 \Rightarrow D = 0.$

Indeed, if $nD = 0$, then we can choose $G = \mathbb{Z}/n$ and the reduced building data $L_i = D, D_g = 0 \forall g \in G$.
 Pardini Equation becomes $nD = 0$, which is satisfied by assumption, so this would define an étale \mathbb{Z}/n -covering $\pi: X \rightarrow Y$ with degree $\deg(\pi) = n$. Thus, π is a topological covering to a simply connected top. space $\Rightarrow \pi$ is an iso, namely $n = \deg(\pi) = 1$.

Let us consider an abelian cover $\pi: X \rightarrow Y$ with reduced building data $\{L_{X_i}, \dots, L_{X_s}\}, \{D_g\}_{g \in G}$.
 Then by Lemma 1: $|X_i| L_{X_i} = \sum_{g \in G} \frac{|X/Y|}{|g|} D_g$, so L_{X_i} is determined by D_g up to torsion.
 Thus, if $\text{Pic}(Y)$ has not torsion (such as for rational varieties $(\mathbb{P}^n, \mathbb{P}^1 \times \mathbb{P}^1, \mathbb{P}^m)$, which are simply connected) then L_{X_i} are uniquely determined by the D_g .
 The only condition to ensure that $\{D_g\}_{g \in G}$ define an abelian covering with group G is that

$\sum_{g \in G} \frac{|x_i| r_{x_i}^g}{|g|} D_g$ is $|x_i|$ -divisible (namely it is linearly equivalent to $|x_i|$ -times another divisor).

In this case, we would define

$$L_{x_i} := \frac{1}{|x_i|} \sum_{g \in G} \frac{|x_i| r_{x_i}^g}{|g|} D_g$$

Then, by the previous Prop, $\{L_{x_i}, \dots, L_{x_s}\}, \{D_g\}_g$ are reduced building data of a unique ab. covering, whose remaining L_x are given by:

$$L_x := \frac{1}{|x|} \sum_{g \in G} \frac{|x| r_x^g}{|g|} D_g, \text{ by Lemma 1.}$$

Def

For this reason, when $\text{Pic}(Y)$ has not torsion, then we say that a collection $\{D_g\}_{g \in G}$ of Y , for which $\sum_{g \in G} \frac{|x_i| r_{x_i}^g}{|g|} D_g$ is $|x_i|$ -divisible, defines a unique abelian cover $\pi: X \rightarrow Y$ with group G .

Example 4


Let $G = \langle e_1, e_2 \rangle \cong \mathbb{Z}/5 \times \mathbb{Z}/5$ and $Y = \mathbb{P}^1 \times \mathbb{P}^1$.

Consider $D_{e_i} = 0 \times \mathbb{P}^1, D_{e_2} = 1 \times \mathbb{P}^1, D_{4e_1+4e_2} = \infty \times \mathbb{P}^1$
 $D_{e_1+2e_2} = \mathbb{P}^1 \times 0, D_{3e_1+4e_2} = \mathbb{P}^1 \times 1, D_{e_1+4e_2} = \mathbb{P}^1 \times \infty$
 $D_g = 0$ otherwise

$\text{Pic}(Y) = \langle H_1, H_2 \rangle_{\mathbb{Z}}$, so let us verify that, given e_1, e_2 dual basis of e_1, e_2 on $G = \langle e_1, e_2 \rangle$, $\sum_{g \in G} \frac{|x_i| r_{x_i}^g}{|g|} D_g$ is

$|\varepsilon_i|$ - divisible by 5:

$\mathbb{P}' \times \mathbb{P}'$



$$\sum_{g \in G} \frac{|\varepsilon_1| r_{\varepsilon_1}^g}{|g|} D_g = \sum_{g \in G} r_{\varepsilon_1}^g D_g = D_{e_1} + 4D_{4e_1+4e_2} + D_{e_1+2e_2} + 3D_{3e_1+4e_2} + D_{e_1+4e_2}.$$

$$\equiv H_1 + 4H_1 + H_2 + 3H_2 + H_2$$

$$\equiv 5(H_1 + H_2) \Rightarrow \mathcal{L}_{\varepsilon_1} := \sigma_{\mathbb{P}' \times \mathbb{P}'}(H_1 + H_2)$$

$$\sum_{g \in G} \frac{|\varepsilon_2| r_{\varepsilon_2}^g}{|g|} D_g = \sum_{g \in G} r_{\varepsilon_2}^g D_g = D_{e_2} + 4D_{4e_1+4e_2} + 2D_{e_1+2e_2} + 4D_{3e_1+4e_2} + 4D_{e_1+4e_2}.$$

$$\equiv H_1 + 4H_1 + 2H_2 + 4H_2 + 4H_2$$

$$\equiv 5(H_1 + 2H_2) \Rightarrow \mathcal{L}_{\varepsilon_2} := \sigma_{\mathbb{P}' \times \mathbb{P}'}(H_1 + 2H_2)$$

Thus $\{D_g\}_{g \in (\mathbb{Z}/5)^2}$ in $\mathbb{P}' \times \mathbb{P}'$ defines a unique abelian covering $\pi: S \rightarrow \mathbb{P}' \times \mathbb{P}'$ with group $G = (\mathbb{Z}/5)^2$.

Def S is called Beauville surface. It is a smooth surface of general type (Kodaira dim. 2)

with $K_S^2 = 8$, $p_g(S) = q(S) = 0$, $\chi(S) = 1$, $e(S) = 4$.

It has been constructed by Beauville in his famous book in an alternative way than how we did (see Exercise X.13(4) in Beauville, Complex Algebraic Surfaces).


It is a very important example in the literature of algebraic surfaces of general type.

Example 5 Let $G = (\mathbb{Z}/n)^k$, $Y = \mathbb{P}^2$, and

let $D_{e_i} := l_1, \dots, D_{e_k} := l_n$, $D_{(n-1)\sum_{i=1}^k e_i} := l_{k+1}$, $D_{\delta_j} = 0$ otherwise
 be $(k+1)$ -lines.

Let $G^* = \langle \varepsilon_1, \dots, \varepsilon_n \rangle$ with ε_i dual character of e_i : $\varepsilon_i(e_j) = e^{\frac{2\pi i}{n} \cdot \delta_{ij}}$.

$$\sum_{\delta \in G} \frac{|\varepsilon_\delta| r_{\varepsilon_\delta}^2}{|\delta|} D_\delta = \sum_{i=1}^n \frac{|\varepsilon_\delta| r_{\varepsilon_\delta}^2}{|e_i|} D_{e_i} + \frac{|\varepsilon_\delta| r_{\varepsilon_\delta}^2}{|(n-1)\sum e_i|} \cdot D_{(n-1)\sum e_i}$$



$$= D_{e_1} + 0 + \dots + 0 + (n-1) D_{(n-1)\sum e_i} \equiv nH$$

$$\Rightarrow L_{\varepsilon_\delta} := \mathcal{O}_{\mathbb{P}^2}(H) \quad \forall \delta = 1, \dots, n$$

Thus $\{D_\delta\}_{\delta \in G}$ defines an abelian covering $\pi: X \rightarrow \mathbb{P}^2$ with group $G = (\mathbb{Z}/n)^k$.

Def $\pi: X \rightarrow \mathbb{P}^2$ is called abelian cover of Hirzebruch

Type. Hirzebruch studied and classified these kind of coverings in "Arrangements of lines and Algebraic Surfaces" (in the sense that he established for any possible configuration of lines what type of surfaces are in Enriques-Kodaira classification). Most of them are smooth, but if we choose special configuration of lines, then they became singular. He studied the slope


$K_S^2/\chi(S)$ of the minimal model S of X
 and found 3 beautiful examples satisfying
 Bogomolov - Miyaoka - Yau equality

$K_S^2 = 9\chi(\mathcal{O}_S)$. It is called **BMY-Inequality**


(They proved VS of gen. type $K_S^2 \leq 9\chi(\mathcal{O}_S)$, Fields Medal 1982)

These examples are obtained with the following configuration of lines:

$G = \mathbb{Z}_5^5$



$G = \mathbb{Z}_3^{11}$
 (Hesse Pencil)



$G = \mathbb{Z}_5^8$

